

## New England Biolabs Certificate of Analysis

**Product Name:** *EnGen Lba Cas12a (Cpf1)*  
**Catalog Number:** *M0653S*  
**Concentration:** *1 µM*  
**Packaging Lot Number:** *10115335*  
**Expiration Date:** *07/2023*  
**Storage Temperature:** *-20°C*  
**Storage Conditions:** *500 mM NaCl, 20 mM Sodium Acetate, 0.1 mM EDTA, 0.1 mM TCEP-HCl, 50% Glycerol, (pH 6.0 @ 25°C)*  
**Specification Version:** *PS-M0653S v2.0*

EnGen Lba Cas12a (Cpf1) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0653SVIAL	ENGEN® LBA CAS12A (CPF1)	10115334	Pass
B6002SVIAL	NEBuffer™ r2.1	10102965	Pass

Assay Name/Specification	Lot # 10115335
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer 2.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Functional Testing (Targeted Digestion)</b> A 20 µl reaction in 1X NEBuffer 2.1 containing 20 nM of 100 bp FAM and ROX-labeled double-stranded target DNA, 100 nM crRNA, and 100 nM EnGen® Lba Cas12a (Cpf1) incubated for 15 minutes at 37°C results in ≥90% targeted digestion of the substrate DNA as determined by capillary electrophoresis.	Pass
<b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of f-300 RNA transcript and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 2.1 containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel	Pass

Assay Name/Specification	Lot # 10115335
<p>electrophoresis.</p> <p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda DNA and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Bhairavi Jani  
Production Scientist  
27 Jul 2021



Mary Neal  
Packaging Quality Control Inspector  
27 Jul 2021