

New England Biolabs Certificate of Analysis

Product Name: Endo Hf
Catalog #: P0703S/L
Concentration: 1,000,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10 µg of denatured RNase B in 1 hour at 37°C in a total reaction volume of 10 µl (10 NEB units = 1 IUB millunit).
Lot #: 0181610
Assay Date: 10/2016
Expiration Date: 10/2018
Storage Temp: -20°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 5 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0703S/L v1.0
Effective Date: 12 Feb 2016

Assay Name/Specification (minimum release criteria)	Lot #0181610
Glycosidase Activity (Endo F2, F3) - A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (PNGase F) - A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-Mannosidase) - A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-Xylosidase) - A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-3 Galactosidase) - A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-4 Galactosidase) - A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass



New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0181610
Glycosidase Activity (β-N-Acetylgalactosaminidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β -N-Acetylgalactosaminidase substrate (GalNAc β 1-4Gal β 1-4Glc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylglucosaminidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β -N-Acetylglucosaminidase substrate (GlcNAc β 1-4GlcNAc β 1-4GlcNAc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-Glucosidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Glucosidase substrate (Glc α 1-6Glc α 1-4Glc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-Neuraminidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Neuraminidase substrate (Neu5Ac α 2-3Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-2 Fucosidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-2Gal β 1-4Glc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Fucosidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Galactosidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-3Gal β 1-4GlcNAc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Mannosidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-3Man β 1-4GlcNAc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-6 Galactosidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-6Gal α 1-6Glc α 1-2Fru-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-6 Mannosidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-6Man α 1-6(Man α 1-3)Man-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0181610
Glycosidase Activity (α-N-Acetylgalactosaminidase) - A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -N-Acetylgalactosaminidase substrate (GalNAc α 1-3(Fuc α 1-2)Gal β 1-4Glc-AMC) and 5,000 units of Endo Hf incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Protease Activity (SDS-PAGE) - A 20 μ l reaction in 1X Glyco Buffer 3 containing 24 μ g of a standard mixture of proteins and a minimum of 5,000 units of Endo Hf incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Protein Purity Assay (SDS-PAGE) - Endo Hf is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass



Authorized by
Derek Robinson
12 Feb 2016



Inspected by
Alicia Bielik
03 Nov 2016

