

## New England Biolabs Certificate of Analysis

**Product Name:** DpnII  
**Catalog Number:** R0543L  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (dam-) in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10140675  
**Expiration Date:** 12/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA  
**Specification Version:** PS-R0543S/L v1.0

DpnII Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0543LVIAL	DpnII	10131718	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10131976	Pass
B0543SVIAL	NEBuffer™ DpnII	10110281	Pass

Assay Name/Specification	Lot # 10140675
<b>Protein Purity Assay (SDS-PAGE)</b> DpnII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer DpnII containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 Units of DpnII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer DpnII containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of DpnII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda dam- DNA with DpnII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	Pass

Assay Name/Specification	Lot # 10140675
fragments, >95% can be recut with DpnII.  <b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer DpnII containing 1 µg of Lambda dam- DNA and a minimum of 100 units of DpnII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.




---

Penghua Zhang  
Production Scientist  
03 Mar 2022




---

Michael Tonello  
Packaging Quality Control Inspector  
03 Mar 2022