

New England Biolabs Certificate of Analysis

Product Name: PvuI-HF[®]
Catalog Number: R3150L
Concentration: 20,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10179437
Expiration Date: 08/2024
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version: PS-R3150S/L/V v2.0

| PvuI-HF [®] Component List | | | |
|-------------------------------------|-------------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R3150LVIAL | PvuI-HF [®] | 10160234 | Pass |
| B7024AVIAL | Gel Loading Dye, Purple (6X) | 10182168 | Pass |
| B6004SVIAL | rCutSmart [™] Buffer | 10182167 | Pass |

| Assay Name/Specification | Lot # 10179437 |
|---|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart [™] Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 200 units of PvuI-HF [®] incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 200 units of PvuI-HF [®] incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart [™] Buffer containing 1 µg of pXba DNA and 1 µl of PvuI-HF [®] incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda-HindIII DNA with PvuI-HF [®] , >95% of the DNA | Pass |


| Assay Name/Specification | Lot # 10179437 |
|---|----------------|
| <p>fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PvuI-HF®.</p> | |
| <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 200 units of PvuI-HF® incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> | Pass |
| <p>Protein Purity Assay (SDS-PAGE) PvuI-HF® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p> | Pass |
| <p>qPCR DNA Contamination (E. coli Genomic) A minimum of 20 units of PvuI-HF® is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p> | Pass |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Jianying Luo
Production Scientist
16 Aug 2022



Michael Tonello
Packaging Quality Control Inspector
28 Apr 2023