

## New England Biolabs Certificate of Analysis

**Product Name:** Standard Taq Reaction Buffer Pack  
**Catalog Number:** B9014S  
**Concentration:** 10 X Concentrate  
**Packaging Lot Number:** 10110280  
**Expiration Date:** 06/2025  
**Storage Temperature:** -20°C  
**Specification Version:** PS-B9014S v2.0  
**Composition (1X):** 10 mM Tris-HCl, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, (pH 8.3 @ 25°C)

Standard Taq Reaction Buffer Pack Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B9021SVIAL	Magnesium Chloride (MgCl <sub>2</sub> ) Solution	10092740	Pass
B9014SVIAL	Standard Taq Reaction Buffer Pack	10102086	Pass

Assay Name/Specification	Lot # 10110280
<b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Standard Taq Reaction Buffer is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>qPCR DNA Contamination (E. coli Genomic, Buffer)</b> A minimum of 1 µl of Standard Taq Reaction Buffer is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
<b>Endonuclease Activity (Nicking, Buffer)</b> A 50 µl reaction in 2X Standard Taq Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 µl reaction in 2X Standard Taq Reaction Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined	Pass


Assay Name/Specification	Lot # 10110280
by agarose gel electrophoresis.	
<p><b>PCR Amplification (5 kb Lambda DNA, Buffer)</b> A 50 µl reaction in Standard Taq Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product.</p>	<b>Pass</b>
<p><b>pH (buffers/solutions)</b> The pH of 10X Standard Taq Reaction Buffer is between pH 8.2 and 8.4 at 25°C.</p>	<b>Pass</b>
<p><b>Phosphatase Activity (pNPP, Buffer)</b> A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl<sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl Standard Taq Reaction Buffer incubated for 4 hours at 37°C yields &lt;0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Christie Vazquez  
Production Scientist  
02 Jun 2021



Michael Tonello  
Packaging Quality Control Inspector  
02 Jun 2021