

New England Biolabs Certificate of Analysis

Product Name: Standard Taq Reaction Buffer Pack
Catalog Number: B9014S
Concentration: 10 X Concentrate
Packaging Lot Number: 10236792
Expiration Date: 08/2028
Storage Temperature: -20°C
Specification Version: PS-B9014S v2.0
Composition (1X): 10 mM Tris-HCl, 50 mM KCl, 1.5 mM MgCl₂, (pH 8.3 @ 25°C)


| Standard Taq Reaction Buffer Pack Component List | | | |
|--|--|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| B9021SVIAL | Magnesium Chloride (MgCl ₂) Solution | 10221498 | Pass |
| B9014SVIAL | Standard Taq Reaction Buffer Pack | 10230998 | Pass |

| Assay Name/Specification | Lot # 10236792 |
|--|----------------|
| Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 2X Standard Taq Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 2X Standard Taq Reaction Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| PCR Amplification (5 kb Lambda DNA, Buffer) A 50 µl reaction in Standard Taq Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product. | Pass |
| Phosphatase Activity (pNPP, Buffer) A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl ₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl Standard Taq Reaction Buffer incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis. | Pass |

| Assay Name/Specification | Lot # 10236792 |
|---|----------------|
| <p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Standard Taq Reaction Buffer is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> | Pass |
| <p>pH (buffers/solutions) The pH of 10X Standard Taq Reaction Buffer is between pH 8.2 and 8.4 at 25°C.</p> | Pass |
| <p>qPCR DNA Contamination (E. coli Genomic, Buffer) A minimum of 1 µl of Standard Taq Reaction Buffer is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p> | Pass |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.


Lea Antonopoulos
Production Scientist
14 Mar 2024


Michael Tonello
Packaging Quality Control Inspector
28 Mar 2024