

New England Biolabs Certificate of Analysis


Product Name: NEB® 5-alpha Competent *E. coli* (Subcloning Efficiency)
Catalog Number: C2988J
Lot Number: 10020866
Expiration Date: 08/2019
Storage Temperature: -80°C
Specification Version: PS-C2988J v1.0

NEB® 5-alpha Competent <i>E. coli</i> (Subcloning Efficiency) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
C2988JVIAL	NEB® 5-alpha Competent <i>E. coli</i> (Subcloning Efficiency)	10009126	Pass


Assay Name/Specification	Lot # 10020866
Antibiotic Sensitivity (Tetracycline) 15 µl of untransformed NEB® 5-alpha Competent <i>E. coli</i> (Subcloning Efficiency) streaked onto a Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.	Pass
Blue-White Screening (α-complementation, Competent Cells) NEB® 5-alpha Competent <i>E. coli</i> (Subcloning Efficiency) were shown to be suitable for blue/white screening by α-complementation of the β-galactosidase gene using pUC19.	Pass
Phage Resistance (φ 80) 15 µl of untransformed NEB® 5-alpha Competent <i>E. coli</i> (Subcloning Efficiency) streaked onto a Rich Broth plate does not support plaque formation by phage φ 80 after incubation for 16 hours at 37°C.	Pass
Transformation Efficiency 50 µl of NEB® 5-alpha Competent <i>E. coli</i> (Subcloning Efficiency) cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10 ⁶ cfu/µg of DNA.	Pass
Antibiotic Sensitivity (Ampicillin) 15 µl of untransformed NEB® 5-alpha Competent <i>E. coli</i> (Subcloning Efficiency) streaked onto a Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.	Pass

Assay Name/Specification	Lot # 10020866
<p>Antibiotic Sensitivity (Chloramphenicol) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Kanamycin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Nitrofurantoin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Nitrofurantoin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Spectinomycin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Streptomycin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Streptomycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Lixin An
Production Scientist
18 May 2018



Corey Rabeau
Packaging Quality Control Inspector
28 Aug 2018