

## New England Biolabs Certificate of Analysis

**Product Name:** T7 Express *lysY* Competent *E. coli* (High Efficiency)  
**Catalog #:** C3010H/I  
**Lot #:** 0071612  
**Assay Date:** 12/2016  
**Expiration Date:** 12/2018  
**Storage Temp:** -80°C  
**Specification Version:** PS-C3010I v1.0  
**Effective Date:** 07 Jul 2016


Assay Name/Specification (minimum release criteria)	Lot #0071612
<b>Antibiotic Resistance (Chloramphenicol)</b> - 15 µl of untransformed T7 Express <i>lysY</i> Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Chloramphenicol will form colonies after incubation for 16 hours at 37°C.	<b>Pass</b>
<b>Antibiotic Resistance (Nitrofurantoin)</b> - 15 µl of untransformed T7 Express <i>lysY</i> Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Nitrofurantoin will form colonies after incubation for 16 hours at 37°C.	<b>Pass</b>
<b>Antibiotic Sensitivity (Ampicillin)</b> - 15 µl of untransformed T7 Express <i>lysY</i> Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.	<b>Pass</b>
<b>Antibiotic Sensitivity (Kanamycin)</b> - 15 µl of untransformed T7 Express <i>lysY</i> Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.	<b>Pass</b>
<b>Antibiotic Sensitivity (Spectinomycin)</b> - 15 µl of untransformed T7 Express <i>lysY</i> Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.	<b>Pass</b>
<b>Antibiotic Sensitivity (Streptomycin)</b> - 15 µl of untransformed T7 Express <i>lysY</i> Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Streptomycin will not form colonies after incubation for 16 hours at 37°C.	<b>Pass</b>
<b>Antibiotic Sensitivity (Tetracycline)</b> - 15 µl of untransformed T7 Express <i>lysY</i> Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.	<b>Pass</b>

## New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0071612
<b>Phage Resistance (Φ 80)</b> - 15 µl of untransformed T7 Express lysY Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate does not support plaque formation by phage Φ 80 after incubation for 16 hours at 37°C.	<b>Pass</b>
<b>Transformation Efficiency</b> - 50 µl of T7 Express lysY Competent <i>E. coli</i> (High Efficiency) cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB -Ampicillin plates at 37°C resulted in >0.6 x 10e9 cfu/µg of DNA.	<b>Pass</b>



Authorized by  
Derek Robinson  
07 Jul 2016



Inspected by  
Lixin An  
07 Dec 2016

