

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: E.coli DNA Ligase

Catalog Number: M0205L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to give 50%

ligation of 6 μg of Lambda-HindIII DNA in 30 minutes at 16°C in a

total reaction volume of 20 μl.

Packaging Lot Number: 10146341
Expiration Date: 03/2024
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 200 µg/ml BSA

, 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0205S/L v2.0

E.coli DNA Ligase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0205LVIAL	E.coli DNA Ligase	10140392	Pass	
B0205SVIAL	E. coli DNA Ligase Reaction Buffer	10102315	Pass	

Assay Name/Specification	Lot # 10146341
Protein Purity Assay (SDS-PAGE) E. coli DNA Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie	Pass
Blue detection.	
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	Pass
and a minimum of 1 µl of E. coli DNA Ligase is incubated at 37°C. After incubation	
for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [3H] E. coli DNA and a minimum of 50 units of E. coli DNA Ligase	
incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of E. coli DNA Ligase incubated for 4 hours at 37°C results in	



M0205L / Lot: 10146341

Page 1 of 2

Assay Name/Specification	Lot # 10146341
<10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of CIP-treated Lambda-HindIII DNA and a minimum of 20 units of E. coli DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of E. coli DNA Ligase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 06 Jun 2022 Erin Varney

Packaging Quality Control Inspector

06 Jun 2022



M0205L / Lot: 10146341