

## New England Biolabs Certificate of Analysis


**Product Name:** M-MuLV Reverse Transcriptase  
**Catalog Number:** M0253L  
**Concentration:** 200,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to incorporate 1 nmol of dTTP into an acid-insoluble form in 10 minutes at 37°C.  
**Packaging Lot Number:** 10061412  
**Expiration Date:** 07/2021  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM Tris-HCl, 150 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 0.1 % IGEPAL® CA-630, 50 % Glycerol, (pH 7.6 @ 25°C)  
**Specification Version:** PS-M0253S/L v1.0

M-MuLV Reverse Transcriptase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0253LVIAL	M-MuLV Reverse Transcriptase	10046979	Pass
B0253SVIAL	M-MuLV Reverse Transcriptase Reaction Buffer	10026291	Pass

Assay Name/Specification	Lot # 10061412
<p><b>RNase Activity Assay (2 Hour Digestion)</b>            A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of M-MuLV Reverse Transcriptase incubated for 2 hours at 37°C results in no detectable degradation of the RNA as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p><b>Endonuclease Activity (Nicking)</b>            A 50 µl reaction in M-MuLV Reverse Transcriptase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 200 units of M-MuLV Reverse Transcriptase incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p><b>Exonuclease Activity (Radioactivity Release)</b>            A 50 µl reaction in M-MuLV Reverse Transcriptase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 200 units of M-MuLV Reverse Transcriptase incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	Pass
<p><b>Non-Specific DNase Activity (16 Hour)</b></p>	Pass

Assay Name/Specification	Lot # 10061412
A 50 µl reaction in M-MuLV Reverse Transcriptase Reaction Buffer containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 200 units of M-MuLV Reverse Transcriptase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

  
\_\_\_\_\_  
David Guo  
Production Scientist  
24 Jul 2019

  
\_\_\_\_\_  
Michael Tonello  
Packaging Quality Control Inspector  
12 Dec 2019