

New England Biolabs Certificate of Analysis

Product Name: *Bst DNA Polymerase, Full Length*
Catalog #: *M0328S/L*
Concentration: *5,000 units/ml*
Unit Definition: *One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 65°C.*
Lot #: *0051509*
Assay Date: *09/2015*
Expiration Date: *09/2017*
Storage Temp: *-20°C*
Storage Conditions: *10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1 % Triton®X-100, 50 % Glycerol, (pH 7.1 @ 25°C)*
Specification Version: *PS-M0328S/L v1.0*
Effective Date: *16 Oct 2015*

| Assay Name/Specification (minimum release criteria) | Lot #0051509 |
|--|--------------|
| <p>Endonuclease Activity (Nicking) - A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of <i>Bst</i> DNA Polymerase, Full Length incubated for 4 hours at either 37°C or 65°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p> | Pass |
| <p>Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 units of <i>Bst</i> DNA Polymerase, Full Length incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> | Pass |
| <p>Phosphatase Activity (pNPP) - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM <i>p</i>-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units <i>Bst</i> DNA Polymerase, Full Length incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p> | Pass |
| <p>Protein Purity Assay (SDS-PAGE) - <i>Bst</i> DNA Polymerase, Full Length is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p> | Pass |
| <p>qPCR DNA Contamination (<i>E. coli</i> Genomic) - A minimum of 5 units of <i>Bst</i> DNA Polymerase, Full Length is screened for the presence of <i>E. coli</i> genomic DNA using SYBR® Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome.</p> | Pass |



**New England Biolabs
Certificate of Analysis**

| Assay Name/Specification (minimum release criteria) | Lot #0051509 |
|---|--------------|
| RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of <i>Bst</i> DNA Polymerase, Full Length is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass |



Authorized by
Melanie Fortier
16 Oct 2015



Inspected by
Katie Gebhardt
24 Nov 2015

