

New England Biolabs Certificate of Analysis

Product Name: ProtoScript[®] II Reverse Transcriptase
Catalog Number: M0368S
Concentration: 200,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme that will incorporate 1 nmol of dTTP into an acid-insoluble form in 10 minutes at 37°C.
Packaging Lot Number: 10198143
Expiration Date: 12/2024
Storage Temperature: -20°C
Storage Conditions: 20 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 0.01 % IGEPAL[®] CA-630, 50 % Glycerol, (pH 7.5 @ 25°C)
Specification Version: PS-M0368S/L/X v2.0


ProtoScript [®] II Reverse Transcriptase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0368SVIAL	ProtoScript [®] II Reverse Transcriptase	10187667	Pass
B1034AVIAL	10X DTT	10170340	Pass
B0368SVIAL	ProtoScript [®] II Reverse Transcriptase Reaction Buffer	10194199	Pass

Assay Name/Specification	Lot # 10198143
<p>Endonuclease Activity (Nicking) A 50 µl reaction in ProtoScript[®] II Reverse Transcriptase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 200 units of ProtoScript[®] II Reverse Transcriptase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in ProtoScript[®] II Reverse Transcriptase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of ProtoScript[®] II Reverse Transcriptase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 200 units of ProtoScript[®] II Reverse Transcriptase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass

Assay Name/Specification	Lot # 10198143
<p>Protein Purity Assay (SDS-PAGE) ProtoScript® II Reverse Transcriptase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity Assay (4 Hour Digestion) A 10 µl reaction in ProtoScript® II Reverse Transcriptase Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of ProtoScript® II Reverse Transcriptase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 200 units of ProtoScript® II Reverse Transcriptase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.


Lea Antonopoulos
Production Scientist
03 Jan 2023


Josh Hersey
Packaging Quality Control Inspector
21 Jul 2023