

## New England Biolabs Certificate of Analysis

**Product Name:** *NudC Pyrophosphatase*  
**Catalog Number:** *M0607S*  
**Concentration:** *10 µM*  
**Unit Definition:** *One µM of NudC Pyrophosphatase can hydrolyze ≥200 µM of NAD<sup>+</sup> into NMN<sup>+</sup> and AMP in 30 minutes at 37°C.*  
**Packaging Lot Number:** *10082274*  
**Expiration Date:** *09/2021*  
**Storage Temperature:** *-20°C*  
**Storage Conditions:** *10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)*  
**Specification Version:** *PS-M0607S v1.0*

NudC Pyrophosphatase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0607SVIAL	NudC Pyrophosphatase	10082275	Pass
B7203SVIAL	NEBuffer™ 3.1	10092685	Pass
B1222AVIAL	0.1M DTT	10065839	Pass

Assay Name/Specification	Lot # 10082274
<b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of NudC Pyrophosphatase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> NudC Pyrophosphatase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 1 µM of NudC Pyrophosphatase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



---

Jenna Ware  
Production Scientist  
02 Feb 2021



---

Michael Tonello  
Packaging Quality Control Inspector  
02 Feb 2021