

## New England Biolabs Certificate of Analysis

**Product Name:** Rapid™ PNGase F  
**Catalog Number:** P0710S  
**Unit Definition:** N/A  
**Packaging Lot Number:** 10162235  
**Expiration Date:** 08/2023  
**Storage Temperature:** 4°C  
**Storage Conditions:** 50 mM NaCl , 20 mM Tris-HCl , 5 mM EDTA, (pH 7.5 @ 25°C)  
**Specification Version:** PS-P0710S v2.0

| Rapid™ PNGase F Component List |                          |            |                      |
|--------------------------------|--------------------------|------------|----------------------|
| NEB Part Number                | Component Description    | Lot Number | Individual QC Result |
| P0710SVIAL                     | Rapid™ PNGase F          | 10156839   | Pass                 |
| B0718SVIAL                     | 5X Rapid PNGase F Buffer | 10156840   | Pass                 |

| Assay Name/Specification   | Lot # 10162235 |
|--|----------------|
| <b>Glycosidase Activity (β-N-Acetylglucosaminidase)</b><br>A 10 µl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 1 µl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass           |
| <b>Glycosidase Activity (β-Xylosidase)</b><br>A 10 µl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 1 µl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.                             | Pass           |
| <b>Glycosidase Activity (β1-4 Galactosidase)</b><br>A 10 µl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC) and 1 µl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.                | Pass           |
| <b>Glycosidase Activity (β-Mannosidase)</b><br>A 10 µl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 1 µl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer  | Pass           |

| Assay Name/Specification   | Lot # 10162235 |
|--|----------------|
| chromatography.  |                |
| <p><b>Glycosidase Activity (<math>\beta</math>-N-Acetylgalactosaminidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\beta</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>                               | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-6 Galactosidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\alpha</math>-Galactosidase substrate (Gal<math>\alpha</math>1-6Gal<math>\alpha</math>1-6Glc<math>\alpha</math>1-2Fru-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>                          | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-6 Mannosidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-6Man<math>\alpha</math>1-6(Man<math>\alpha</math>1-3)Man-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>                            | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>-Glucosidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\alpha</math>-Glucosidase substrate (Glc<math>\alpha</math>1-6Glc<math>\alpha</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>  | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>-N-Acetylgalactosaminidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\alpha</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\alpha</math>1-3(Fuc<math>\alpha</math>1-2)Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>-Neuraminidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\alpha</math>-Neuraminidase substrate (Neu5Ac<math>\alpha</math>2-3Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\beta</math>1-3 Galactosidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\beta</math>-Galactosidase substrate (Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>                            | <b>Pass</b>    |

| Assay Name/Specification   | Lot # 10162235 |
|--|----------------|
| <p><b>Glycosidase Activity (<math>\alpha</math>1-3 Fucosidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\alpha</math>-Fucosidase substrate (Fuca1-3Gal<math>\beta</math>1-4GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-3 Galactosidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\alpha</math>-Galactosidase substrate (Gal<math>\alpha</math>1-3Gal<math>\beta</math>1-4GlcNAc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>                         | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-3 Mannosidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-3Man<math>\beta</math>1-4GlcNAc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>                             | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-2 Fucosidase)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled <math>\alpha</math>-Fucosidase substrate (Fuca1-2Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>  | <b>Pass</b>    |
| <p><b>Protease Activity (SDS-PAGE)</b><br/>A 20 <math>\mu</math>l reaction in 1X Rapid PNGase F Buffer containing 24 <math>\mu</math>g of a standard mixture of proteins and a minimum of 5 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>   | <b>Pass</b>    |
| <p><b>Protein Purity Assay (SDS-PAGE)</b><br/>Rapid PNGase F is <math>\geq</math> 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>  | <b>Pass</b>    |
| <p><b>Glycosidase Activity (Endo F1, F2, H)</b><br/>A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 1 <math>\mu</math>l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>  | <b>Pass</b>    |
| <p><b>Glycosidase Activity (Endo F2, F3)</b></p>   | <b>Pass</b>    |

| Assay Name/Specification   | Lot # 10162235 |
|--|----------------|
| A 10 µl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 1 µl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Alicia Bielik  
Production Scientist  
08 Sep 2022



Michael Tonello  
Packaging Quality Control Inspector  
08 Sep 2022