

New England Biolabs Certificate of Analysis

Product Name: *Endoglycoceramidase I (EGCase I)*
Catalog Number: P0773S
Concentration: 6 U/ml
Unit Definition: One unit of *R. triatomea* EGCase I is defined as the amount of enzyme required to hydrolyze 1 μ mol of ganglioside GM1a per minute at 37°C.
Packaging Lot Number: 10056389
Expiration Date: 12/2021
Storage Temperature: -20°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0773S v1.0

| Endoglycoceramidase I (EGCase I) Component List | | | |
|---|----------------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| P0773SVIAL | Endoglycoceramidase I (EGCase I) | 10057247 | Pass |
| B0773SVIAL | EGCase I Buffer | 10031041 | Pass |

| Assay Name/Specification | Lot # 10056389 |
|---|----------------|
| <p>Glycosidase Activity (Endo F1, F2, H) A 10 μl reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (Endo F2, F3) A 10 μl reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (PNGase F) A 10 μl reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α-Glucosidase) A 10 μl reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled</p> | Pass |

| Assay Name/Specification | Lot # 10056389 |
|---|----------------|
| <p>α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | |
| <p>Glycosidase Activity (α-Neuraminidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α1-2 Fucosidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-2Galβ1-4Glc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α1-3 Fucosidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α1-3 Galactosidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α1-3 Mannosidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-3Manβ1-4GlcNAc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α1-6 Galactosidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α1-6 Mannosidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 6 mU of EGCCase I</p> | Pass |

| Assay Name/Specification | Lot # 10056389 |
|---|----------------|
| <p>incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | |
| <p>Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fuca1-2)Galβ1-4Glc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (β-Mannosidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (β-Xylosidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μl reaction in EGCCase I Buffer containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 6 mU of EGCCase I incubated for 20 hours at 37°C results in no detectable activity as</p> | Pass |

