

## New England Biolabs Certificate of Analysis


**Product Name:** BsmI  
**Catalog #:** R0134S/L  
**Concentration:** 10,000 units/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 65°C in a total reaction volume of 50 µl.  
**Lot #:** 0161608  
**Assay Date:** 08/2016  
**Expiration Date:** 8/2018  
**Storage Temp:** -20°C  
**Storage Conditions:** 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
**Specification Version:** PS-R0134S/L v1.0  
**Effective Date:** 29 Jul 2013

| Assay Name/Specification (minimum release criteria)  | Lot #0161608 |
|--|--------------|
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 100 units of BsmI incubated for 4 hours at 65°C releases <0.1% of the total radioactivity. | <b>Pass</b>  |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda DNA with BsmI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated fragments, >95% can be recut with BsmI.  | <b>Pass</b>  |
| <b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of 30 units of BsmI incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.      | <b>Pass</b>  |

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by  
Derek Robinson  
29 Jul 2013



Inspected by  
Penghua Zhang  
05 Aug 2016

