

New England Biolabs Certificate of Analysis

Product Name: *ApaLI*
Catalog Number: *R0507M*
Concentration: *50,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10232749*
Expiration Date: *10/2025*
Storage Temperature: *-20°C*
Storage Conditions: *10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)*
Specification Version: *PS-R0507T/M v2.0*


| ApaLI Component List | | | |
|----------------------|-----------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R0507MVIAL | ApaLI | 10210765 | Pass |
| B6004SVIAL | rCutSmart™ Buffer | 10228580 | Pass |

| Assay Name/Specification | Lot # 10232749 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled M13mp18 DNA and a minimum of 50 units of ApaLI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of ApaLI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and 1 µl of ApaLI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with ApaLI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, | Pass |

| Assay Name/Specification | Lot # 10232749 |
|---|--------------------|
| <p>>95% can be recut with ApaLI.</p> <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 100 units of ApaLI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> | <p>Pass</p> |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



YunJie Sun
Production Scientist
10 Oct 2023



Michael Tonello
Packaging Quality Control Inspector
20 Mar 2024