

New England Biolabs Certificate of Analysis

Product Name: *BsII*
Catalog Number: *R0555S*
Concentration: *10,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 55°C in a total reaction volume of 50 µl.*
Lot Number: *10027308*
Expiration Date: *11/2020*
Storage Temperature: *-20°C*
Storage Conditions: *50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA*
Specification Version: *PS-R0555S/L v1.0*


| BsII Component List | | | |
|---------------------|-----------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R0555SVIAL | BsII | 10027307 | Pass |
| B7204SVIAL | CutSmart® Buffer | 10021118 | Pass |

| Assay Name/Specification | Lot # 10027308 |
|---|----------------|
| <p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of BsII incubated for 4 hours at 55°C releases <0.1% of the total radioactivity.</p> | Pass |
| <p>Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BsII, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BsII.</p> | Pass |
| <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 30 units of BsII incubated for 16 hours at 55°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> | Pass |

This product has been tested and shown to be in compliance with all specifications.



Tony Spear-Alfonso
Production Scientist
01 Oct 2018



Michael Tonello
Packaging Quality Control Inspector
12 Nov 2018