240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: SrfI

Catalog #: R0629S/L
Concentration: 20,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pNEB193-Srf1 DNA in CutSmart incubated for 1 hour

at 37°C in a total reaction volume of 50 μ l.

 Lot #:
 0011603

 Assay Date:
 03/2016

 Expiration Date:
 9/2017

 Storage Temp:
 -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-R0629S/L v1.0
Effective Date: 11 Nov 2015

Assay Name/Specification (minimum release criteria)	Lot #0011603
Endonuclease Activity (Nicking) - A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled pBR322 DNA and a minimum of 100 units of Srf1 incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 200 units of Srf1 incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) - A 50 μl reaction in CutSmart® Buffer containing 1 μg of pNEB193-SrfI DNA and 1 μl of SrfI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) - After a 20-fold over-digestion of pNEB193-Srfl DNA with Srfl, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Srfl.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart® Buffer containing 1 µg of pNEB193-Srfl DNA and a minimum of 20 units of Srfl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) - Srfl is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

Authorized by Derek Robinson 11 Nov 2015







Inspected by Penghua Zhang 01 Mar 2016

holog