

## New England Biolabs Certificate of Analysis

**Product Name:** Control LAMP Primer Mix (rActin)  
**Catalog Number:** S0164S  
**Concentration:** 10 X Concentrate  
**Packaging Lot Number:** 10207319  
**Expiration Date:** 08/2025  
**Storage Temperature:** -20°C  
**Specification Version:** PS-S0164S v1.0  
**Composition (1X):** Proprietary


Control LAMP Primer Mix (rActin) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
S0164AVIAL	Internal Control LAMP Primer Mix (rActin)	10202231	Pass

Assay Name/Specification	Lot # 10207319
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 µl of Control LAMP Primer Mix (rActin) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Functional Testing (LAMP primers)</b> A 25 µl reaction in 1X WarmStart® LAMP Master Mix with UDG in the presence of LAMP Fluorescent Dye and 1X Control LAMP Primer Mix (rActin) containing 10 ng human RNA results in a threshold time of ≤ 20 minutes as determined by fluorescent detection. Reactions that lack human RNA template remain negative over a 30 minute incubation at 65°C.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 µl of Control LAMP Primer Mix (rActin) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Phosphatase Activity (pNPP)</b> A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl <sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 20 µl of Control LAMP Primer Mix (rActin) incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase	Pass

Assay Name/Specification	Lot # 10207319
<p>activity as determined by spectrophotometric analysis.</p> <p><b>RNase Activity (Extended Digestion)</b>            A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Control LAMP Primer Mix (rActin) is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.

  
 Lea Antonopoulos  
 Production Scientist  
 23 Aug 2023

  
 Josh Hersey  
 Packaging Quality Control Inspector  
 25 Oct 2023