

## New England Biolabs Product Specification

<i>Product Name:</i>	<i>Rapid™ PNGase F</i>
<i>Catalog #:</i>	<i>P0710S</i>
<i>Concentration:</i>	<i>1 reaction/μl</i>
<i>Unit Definition:</i>	<i>N/A</i>
<i>Shelf Life:</i>	<i>12 months</i>
<i>Storage Temp:</i>	<i>4°C</i>
<i>Storage Conditions:</i>	<i>50 mM NaCl, 20 mM Tris-HCl, 5 mM EDTA, (pH 7.5 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-P0710S v2.0</i>
<i>Effective Date:</i>	<i>16 Jan 2018</i>

### Assay Name/Specification (minimum release criteria)

**Glycosidase Activity (Endo F1, F2, H)** - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity (Endo F2, F3)** - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity (β-Mannosidase)** - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity (β-N-Acetylgalactosaminidase)** - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity (β-N-Acetylglucosaminidase)** - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity (β-Xylosidase)** - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.



## New England Biolabs Product Specification

### Assay Name/Specification (minimum release criteria)

**Glycosidase Activity ( $\beta$ 1-3 Galactosidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\beta$ -Galactosidase substrate (Gal $\beta$ 1-3GlcNAc $\beta$ 1-4Gal $\beta$ 1-4Glc-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity ( $\beta$ 1-4 Galactosidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\beta$ -Galactosidase substrate (Gal $\beta$ 1-4GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity ( $\alpha$ -Glucosidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\alpha$ -Glucosidase substrate (Glc $\alpha$ 1-6Glc $\alpha$ 1-4Glc-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity ( $\alpha$ -N-Acetylgalactosaminidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\alpha$ -N-Acetylgalactosaminidase substrate (GalNAc $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal $\beta$ 1-4Glc-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity ( $\alpha$ -Neuraminidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\alpha$ -Neuraminidase substrate (Neu5Ac $\alpha$ 2-3Gal $\beta$ 1-3GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity ( $\alpha$ 1-2 Fucosidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\alpha$ -Fucosidase substrate (Fuc $\alpha$ 1-2Gal $\beta$ 1-4Glc-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity ( $\alpha$ 1-3 Fucosidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\alpha$ -Fucosidase substrate (Fuc $\alpha$ 1-3Gal $\beta$ 1-4GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity ( $\alpha$ 1-3 Galactosidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\alpha$ -Galactosidase substrate (Gal $\alpha$ 1-3Gal $\beta$ 1-4GlcNAc-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity ( $\alpha$ 1-3 Mannosidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\alpha$ -Mannosidase substrate (Man $\alpha$ 1-3Man $\beta$ 1-4GlcNAc-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Glycosidase Activity ( $\alpha$ 1-6 Galactosidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\alpha$ -Galactosidase substrate (Gal $\alpha$ 1-6Gal $\alpha$ 1-6Glc $\alpha$ 1-2Fru-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.



## New England Biolabs Product Specification

**Assay Name/Specification (minimum release criteria)**

**Glycosidase Activity ( $\alpha$ 1-6 Mannosidase)** - A 10  $\mu$ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled  $\alpha$ -Mannosidase substrate (Man $\alpha$ 1-6Man $\alpha$ 1-6(Man $\alpha$ 1-3)Man-AMC) and 1  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.

**Protease Activity (SDS-PAGE)** - A 20  $\mu$ l reaction in 1X Rapid PNGase F Buffer containing 24  $\mu$ g of a standard mixture of proteins and a minimum of 5  $\mu$ l of Rapid PNGase F incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.

**Protein Purity Assay (SDS-PAGE)** - Rapid PNGase F is  $\geq$  95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.



Date 16 Jan 2018

Derek Robinson  
Director of Quality Control

