

New England Biolabs Product Specification

<i>Product Name:</i>	<i>SmaI</i>
<i>Catalog #:</i>	<i>R0141S/L/E</i>
<i>Concentration:</i>	<i>20,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml rAlbumin (pH 7.4 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-R0141S/L/E v2.0</i>
<i>Effective Date:</i>	<i>04 Aug 2022</i>

Assay Name/Specification (minimum release criteria)

Blue-White Screening (Terminal Integrity) - A sample of pUC19 vector linearized with a 10-fold excess of SmaI, religated and transformed into an *E. coli* strain expressing the LacZ beta fragment gene results in <1% white colonies.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of Lambda DNA with SmaI, ~50% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SmaI.

Endonuclease Activity (Nicking) - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of SmaI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 100 units of SmaI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Functional Testing (15 minute Digest) - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and 1 µl of SmaI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 60 units of SmaI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - SmaI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.



New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)

qPCR DNA Contamination (<i>E. coli</i> Genomic) - A minimum of 20 units of SmaI is screened for the presence of <i>E. coli</i> genomic DNA using SYBR® Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome.

One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.



Date 04 Aug 2022

Derek Robinson
Quality Approver

