

New England Biolabs Product Specification

Product Name:	<i>NotI</i>
Catalog #:	R0189M
Concentration:	50,000 units/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 µg of pBC4 DNA in NEBuffer r3.1 in 1 hour at 37°C in a total reaction volume of 50 µl.
Shelf Life:	24 months
Storage Temp:	-20°C
Storage Conditions:	10 mM Tris-HCl, 250 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton X-100, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version:	PS-R0189M v2.0
Effective Date:	17 Mar 2024

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of NotI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 100 units of NotI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of pBC4 DNA with NotI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NotI.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of pBC4 DNA and a minimum of 100 units of NotI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - NotI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

qPCR DNA Contamination (*E. coli* Genomic) - A minimum of 10 units of NotI is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.





240 County Road
Ipswich, MA 01938-2723

Tel 978-927-5054
Fax 978-921-1350

www.neb.com
info@neb.com

New England Biolabs Product Specification

*One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.*

Nancy Considine

Date 17 Mar 2024

Nancy Considine
Quality Approver

