

Lambda

48,502 base pairs

GenBank Accession #: NC_001416

See page 128 for ordering information.

There are no restriction sites for the following enzymes: AbsI(x), AsiSI, FseI, MauBI(x), MreI(x), NotI, PacI, SfiI, SpeI, SrfI, SwaI

(x) = enzyme not available from NEB




We recommend NEBcutter at NEBcutter.com to identify the restriction sites within your DNA sequence. NEBcutter indicates cut frequency and methylation-state sensitivity.

Lambda (λ) is a large, temperate *E. coli* bacteriophage with a linear, largely double-stranded DNA genome (1-5). At each end, the 5' strand overhangs the 3' strand by 12 bases. These single-stranded overhangs are complementary and anneal to form a *cos* site following entry into a host cell. Once annealed, the genome is a circular, completely double-stranded molecule which serves as a template for rolling-circle replication.

Many laboratory strains of lambda are derivatives of the strain λ *cl857 ind1 Sam7*, which contains four point mutations relative to the wild type strain. The *ind1* mutation in the *ci* gene creates a new HindIII site at 37584 not present in the wild type. All lambda products sold by NEB are λ *cl857 ind1 Sam7*.

Numbering of the genome sequence begins at the first (5'-most) base of the left end (bottom of diagram below) and continues rightward from late genes *nu1* and *A* towards the early genes. The map below shows the positions of all known ORFs larger than 200 codons.

Enzymes with unique restriction sites are shown in **bold** type and enzymes with two restriction sites are shown in regular type. **Coordinates indicate position of cutsite on the top strand. In previous catalogs, coordinates referred to the position of the 5' most base on the top strand, please make note of new numbering system.**



References

- (1) Echols, H. and Murialdo, H. (1978) *Microbiol. Rev.*, 42, 577-591.
- (2) Szybalski, E.H. and Szybalski, W. (1979) *Gene*, 7, 217-270
- (3) Daniels, D.L., de Wet, J.R. and Blattner, F.R. (1980) *J. Virol.*, 33, 390-400.
- (4) Sanger, F., Coulson, A.R., Hong, G.F., Hill, D.F. and Petersen, G.B. (1982) *J. Mol. Biol.*, 162, 729-773.
- (5) Daniels, D.L. et al. (1983). In R.W. Hendrix, J.W. Roberts, F.W. Stahl and R.A. Weisberg (Eds.), *Lambda II: Appendix*, New York: Cold Spring Harbor Press.